

A higher level of human papillomavirus 16 DNA was associated with an increased risk for cervical carcinoma in situ

Josefsson AM, Magnusson PK, Ylitalo N, et al. Viral load of human papilloma virus 16 as a determinant for development of cervical carcinoma in situ: a nested case-control study. *Lancet*. 2000 Jun 24;355:2189-93.

QUESTION

In women with a first normal cervical smear during screening, what is the association between human papillomavirus (HPV) DNA levels and development of cervical carcinoma in situ (CIS)?

DESIGN

Nested case-control study of women in a screening program from 1969 to 1995.

SETTING

Uppsala County, Sweden.

PARTICIPANTS

Women who were < 50 years of age at entry, were born in Sweden, and had a normal first cervical smear and smears that contained genomic DNA as measured by β -actin. The case group consisted of 478 women (2081 smears). 5 women in the control group were matched to each woman with CIS by date of entry to the cohort, age, and time of first smear. Women in the control group had to be alive without developing CIS or invasive cervical cancer or having a hysterectomy before the date of diagnosis for their matched case; when only 1 smear was available for a woman randomly selected from the original pool of women in the control group, a second matched woman from the control group was randomly selected ($n = 608$ [478 first controls and 130 second controls], 1754 smears).

ASSESSMENT OF RISK FACTORS

The level of HPV-16 DNA was estimated by using quantitative polymerase chain reaction (5' exonuclease [Taqman] method). To adjust for different levels of genomic DNA, a nuclear gene (β -actin) was assessed.

MAIN OUTCOME MEASURES

Data on women with CIS were obtained from the National Cancer Registry. An experienced pathologist reassessed the histologic samples to confirm the diagnosis.

MAIN RESULTS

871 smears (42%) from women with CIS and 117 smears (7%) from the control group were positive for HPV-16. After adjustment for level of β -actin, a greater mean level of HPV-16 DNA was associ-

ated with an increased risk for CIS (Table). A similar relation existed when only the first smear was included in the analysis.

CONCLUSION

In women who had a first normal cervical smear in a screening program, a higher mean level of HPV-16 DNA was associated with an increased risk for cervical carcinoma in situ.

Sources of funding: National Institutes of Health; Swedish National Cancer Society; Danish National Science Foundation.

For correspondence: Professor U.B. Gyllensten, Department of Genetics and Pathology, Unit of Medical Genetics, Rudbeck Laboratory, University of Uppsala, SE-751 85 Uppsala, Sweden. FAX 46-18-471-4931. ■

Associations between levels of human papillomavirus (HPV)-16 DNA and risk for cervical cancer

Quintiles of mean HPV-16 threshold cycle number(Ct)*	Adjusted odds ratio (95% CI)†
HPV-16 negative	1.0
HPV-16 positive, Ct value 44.8 to 50.0	2.0 (1.1 to 3.8)
HPV-16 positive, Ct value 41.25 to 44.8	4.4 (2.3 to 8.3)
HPV-16 positive, Ct value 38.99 to 41.25	8.1 (3.8 to 17.3)
HPV-16 positive, Ct value 36.66 to 38.99	18.7 (7.1 to 49.5)
HPV-16 positive, Ct value <36.66	68.8 (15.8 to 299.6)

*Lower values of Ct denote higher levels of HPV-16 DNA.

†Odds ratio adjusted for level of β -actin.

COMMENTARY

Infection with HPV type 16 or 18 has been associated with a higher rate of progression of cervical squamous intraepithelial lesions (SILs) and cancer (1). The ability to identify patients with oncogenic HPV types might lead to improved detection of women who are more likely to develop high-grade cervical SILs and cancer. This improved detection could affect management decisions for women with low-grade cytologic abnormalities. Currently available commercial tests identify the presence of high-risk HPV types. Despite the potential value of this technology, recent reports suggest a lack of utility for these qualitative tests in women with low-grade cervical SILs. The Atypical Squamous Cells of Undetermined Significance/Low-Grade Squamous Epithelial Lesions Triage Study (ALTS) is a large randomized trial of women with recently detected atypical squamous cells of undetermined significance and low-grade cervical SILs. High-risk HPV DNA was detected in 82.9% of women with low-grade cervical SILs, therefore limiting the usefulness of measuring it. Results of the ALTS study for women with atypical squamous cells of undetermined

significance are pending (2). Manos and colleagues studied HPV testing combined with ThinPrep in women with atypical squamous cells of undetermined significance and found that the HPV test had a sensitivity of 89.2% and a specificity of 64.1% (3). Other studies have shown sensitivities $\geq 90\%$ for the second-generation, qualitative, high-risk HPV test (4); however, its false-positive rate ranged from 5% to 20%.

Josefsson and colleagues did a nested case-control study on cervical smears of women who developed cervical CIS and those who did not. They measured HPV-16 viral load on archived cervical smears using a quantitative polymerase chain reaction assay available only in research laboratories. They found a strong positive relation between viral load and risk for cervical CIS, but only 283 of 495 women with CIS had any HPV-16 DNA; of those, only 110 women had viral loads in the top 2 quintiles. In the second study based on the same cohort, the time from high HPV-16 viral load to appearance of CIS was determined: a mean time of 17 years.

(continued on page 35)