

# Review: Antidepressants increase remission and clinical improvement in bulimia nervosa

Bacaltchuk J, Hay P. Antidepressants versus placebo for people with bulimia nervosa. *Cochrane Database Syst Rev.* 2001;(4):CD003931 (latest version 19 Aug 2001).

## QUESTION

In patients with bulimia nervosa (BN), are antidepressants effective for increasing remission and clinical improvement?

## DATA SOURCES

Studies were identified by searching MEDLINE; EMBASE/Excerpta Medica; LILACS; PsycLIT; SCISEARCH; the Cochrane Depression, Anxiety, and Neurosis Group Database of Trials; the Cochrane Controlled Trials Register; *Clinical Evidence*; and reference lists. The *International Journal of Eating Disorders* was also hand searched, and authors and pharmaceutical companies were contacted.

## STUDY SELECTION

Studies were selected if they were randomized controlled trials (RCTs) that compared any antidepressant with placebo for  $\geq 4$  weeks in patients with BN. RCTs were excluded if patients had binge-eating or purging-type anorexia nervosa or binge-eating disorder.

## DATA EXTRACTION

2 reviewers assessed the quality of RCTs and extracted data on patients, study characteristics, drug regimens, and outcomes (including remission [100% reduction in binge or purge episodes], clinical improvement [ $\geq 50\%$  reduction in binge or purge episodes], and dropouts).

## MAIN RESULTS

16 RCTs (1300 patients) met the selection criteria. Any antidepressant was better than placebo for increasing remission at a mean follow-up of 8 weeks (8 RCTs) and clinical improvement at a mean follow-up of 9 weeks (8 RCTs) (Table). Groups did not differ for dropout rates (14 RCTs) (Table).

## CONCLUSION

In patients with bulimia nervosa, antidepressants are effective in the short term for increasing remission and clinical improvement rates.

*Source of funding:* Not stated.

*For correspondence:* Dr. J. Bacaltchuk, Universidade Federal de São Paulo, São Paulo—SP, Brazil. E-mail [bacaltc@ibm.net](mailto:bacaltc@ibm.net). ■

### Antidepressants vs placebo for bulimia nervosa at 6 to 16 weeks\*

Outcomes	Antidepressant type	Weighted event rates	RBI (95% CI)	NNT (CI)
Remission	All types	20% vs 7.9%	105% (32 to 219)	9 (6 to 16)
	TCA	14% vs 9.1%	136% (-4 to 476)	Not significant
	MAOI	25% vs 6.3%	229% (-22 to 1289)	Not significant
	Other antidepressants	15% vs 0%	664% (-1 to 5773)	7 (4 to 27)
Clinical improvement	All types	64% vs 33%	84% (38 to 145)	4 (3 to 5)
	TCA	77% vs 17%	294% (59 to 872)	2 (2 to 3)
	SSRI	58% vs 38%	51% (26 to 81)	5 (4 to 9)
	Other antidepressants	44% vs 8.2%	321% (74 to 919)	3 (3 to 5)
Dropouts	SSRI	34% vs 40%	18% (1 to 32)	Not significant
	Other antidepressants	30% vs 32%	6% (-83 to 52)	Not significant
			RRR (CI)	NNT
	All types	34% vs 31%	3% (-20 to 32)	Not significant
	TCA	26% vs 11%	93% (15 to 225)	Not significant
	MAOI	34% vs 29%	20% (-33 to 113)	Not significant

\*MAOIs = monoamine oxidase inhibitors; SSRIs = selective serotonin reuptake inhibitors; TCAs = tricyclic antidepressants. Other abbreviations defined in Glossary; RBI, RRR, RRI, NNT, NNH, and CI calculated from data in article.

## COMMENTARY

The reviews by Bacaltchuk and colleagues are laudable for the rigor of the data analyses, but they rightly generate more questions than answers. Bacaltchuk and Hay have comprehensively reviewed 16 published RCTs of antidepressant treatments for BN. Although modest effectiveness is shown, high dropout rates among patients limit the clinical application of these data, and the authors comment on the need for more studies of tolerability and cost-effectiveness. The studies included were generally of short duration in young adult women who did not have any substantial psychiatric comorbid conditions. The results therefore cannot be generalized to the substantial minority of bulimic patients with comorbid "multi-impulsive" personality characteristics (1) or substance abuse or to adolescents.

Pharmacologic treatment trials of BN are dominated by the reported reduction in bulimic symptoms, but clinicians and their patients are more interested in remission of symptoms. The emphasis of this review on remission is therefore of greater clinical application than the emphasis of its sources. The review discusses the limitations of outcome measures and is right to conclude that the use of antidepressants as sole therapy "does not seem sufficient to effectively treat these patients."

Bacaltchuk and colleagues review a scant number of studies comparing combined antidepressant medication and psychotherapy with each

treatment alone. In clinical practice, cognitive behavioral therapy (CBT), which is limited by its availability, is generally regarded as the treatment of choice for BN, with antidepressant medication as an adjunct. This review supports that approach by using restricted data from fairly small studies. However, the clinical risk associated with a pharmacologic approach to BN seems to be a higher dropout rate than with CBT, and again, the results cannot necessarily be generalized beyond young adult women who have no substantial comorbid illness.

The U.K. Department of Health's National Service Framework for Mental Health has stressed the importance of managing such eating disorders as BN in primary care (2), noting that "antidepressants can reduce purging and bingeing whether or not the person is also depressed." Although this statement is true in the short term, it would seem an optimistic reading of the literature. Prescription of antidepressants may appear to be the easiest route in a primary care setting, but the clinical implication of Bacaltchuk and colleagues' review is that the easiest route may not be the most effective, cost-effective, or acceptable for clinicians and their patients.

However, in the busy world of primary care, the treatment of BN will continue to be driven by available resources. CBT for BN is generally preferred by the family doctor when specialists with such training

(continued on page 107)

# Review: Psychological treatment is as effective as antidepressants for bulimia nervosa, but a combination is best

Bacaltchuk J, Hay P, Trefiglio R. Antidepressants versus psychological treatments and their combination for bulimia nervosa. *Cochrane Database Syst Rev.* 2001;(4):CD003385 (latest version 13 Aug 2001).

## QUESTIONS

In patients with bulimia nervosa (BN), are antidepressants as effective as psychological treatment (PT) for increasing remission and clinical improvement rates? Is a combination of antidepressants and PT better than each intervention alone?

## DATA SOURCES

Studies were identified by searching MEDLINE; EMBASE/Excerpta Medica; LILACS; PsycLIT; SCISEARCH; the Cochrane Depression, Anxiety, and Neurosis Group Database of Trials; the Cochrane Controlled Trials Register; *Clinical Evidence*; and reference lists. The *International Journal of Eating Disorders* and book chapters on BN were also hand searched, and authors and pharmaceutical companies were contacted.

## STUDY SELECTION

Studies were selected if they were randomized controlled trials (RCTs) that compared antidepressants with PT in patients with BN. Studies were excluded if patients had binge-eating or purging-type anorexia nervosa or binge-eating disorder.

## DATA EXTRACTION

2 reviewers assessed the quality of studies and extracted data on patients, study characteristics, interventions, and outcomes (including remission [100% reduction in binge or purge episodes], clinical improvement [ $\geq 50\%$  reduction in binge or purge episodes], and dropouts).

## MAIN RESULTS

5 RCTs (237 patients) compared antidepressants with PT. Groups did not differ significantly for remission (5 RCTs); only 1 RCT reported on clinical improvement. More dropouts occurred in the antidepressant group than in the PT group (4 RCTs) (Table). 5 RCTs (247 patients) compared combination and single interventions.

*Antidepressants vs combination:* More patients in the combination group than in the antidepressant-alone group had remission (4 RCTs) (Table); only 1 RCT reported on clinical improvement. Groups did not differ for dropout rates (4 RCTs).

*PT vs combination:* More patients in the combination group than in the PT-alone

group had remission (6 RCTs); fewer patients in the PT-alone group than in the combination group dropped out (6 RCTs) (Table). Groups did not differ for clinical improvement (2 RCTs) (Table).

## CONCLUSIONS

In patients with bulimia nervosa, psychological treatment (PT) and antidepressants do not differ in remission rates, but dropout rates are lower with PT. A combination of antidepressants and PT is best for increasing remission.

*Source of funding:* Not stated.

*For correspondence:* Dr. J. Bacaltchuk, Universidade Federal de São Paulo, São Paulo—SP, Brazil. E-mail [bacaltc@ibm.net](mailto:bacaltc@ibm.net). ■

## Antidepressants (AD) vs psychological treatment (PT) for bulimia nervosa\*

Outcomes	Comparisons	Weighted event rates	RBI (95% CI)	NNT (CI)
Remission	PT vs AD	41% vs 20%	63% (−14 to 210)	Not significant
	AD + PT vs AD	47% vs 23%	79% (11 to 188)	5 (3 to 21)
	PT + AD vs PT	50% vs 36%	30% (1 to 68)	8 (5 to 37)
			<b>RBR (CI)</b>	<b>NNH</b>
Clinical improvement	PT + AD vs PT	46% vs 52%	8% (−70 to 50)	Not significant
			<b>RRR (CI)</b>	<b>NNT (CI)</b>
Dropouts	PT vs AD	18% vs 41%	54% (9 to 76)	5 (3 to 10)
	AD + PT vs AD	35% vs 41%	16% (−45 to 51)	Not significant
			<b>RRI (CI)</b>	<b>NNH (CI)</b>
	PT + AD vs PT	26% vs 16%	74% (14 to 167)	10 (6 to 40)

\*RBR = relative benefit reduction. Other abbreviations defined in Glossary; RBI, RBR, RRR, RRI, NNT, NNH, and CI calculated from data in article. Follow-up ranged from 5 to 24 weeks.

## COMMENTARY (continued from page 106)

are available. But the Royal College of Psychiatrists, in collaboration with the Consumers' Association, has recently reported the dearth of specialist eating-disorder services beyond southeastern England (3). Thus, in the more likely scenario of limited eating-disorder services, use of antidepressant medication may seem more attractive. These 2 reviews agree with that approach and suggest that antidepressant medication will produce positive short-term results. BN, however, is not a short-term illness. Relapse prevention deserves greater scrutiny for patients with BN and anorexia nervosa, and longer-term follow-up studies should drive the next generation of treatment intervention studies.

Regarding treatment of BN in particular, a pressing need exists for longer-term studies examining relapse rates, health economics, and comparisons of classes of antidepressants for treatment concordance.

John F Morgan, MD, MA  
St. George's Hospital Medical School, University of London  
London, England, UK

## References

- Lacey JH. Inpatient treatment of multi-impulsive bulimia nervosa. In: Brownell KD, Fairburn CG, eds. *Eating Disorders and Obesity: A Comprehensive Handbook*. New York: Guilford Press;1995:361-8.
- A National Service Framework for Mental Health Modern Standards and Service Models. London: Department of Health;1999.
- Eating Disorders in the UK: Policies for Service Development and Training. London: Royal College of Psychiatrists; August 2001. <http://www.rcpsych.ac.uk/publications/cr/council/cr87.pdf>.