

# Review: Evidence on surgical interventions for distal radial fractures is inconclusive

Handoll HH, Madhok R. *Surgical interventions for treating distal radial fractures in adults.* Cochrane Database Syst Rev. 2001(4):CD003209 (latest version 29 Mar 2001).

## QUESTION

In adults with fractures of the distal radius, is surgical treatment effective for improving clinical outcome?

## DATA SOURCES

Studies were identified by searching 6 databases; hand searching conference proceedings; and scanning reference lists.

## STUDY SELECTION

Randomized or quasirandomized controlled trials were selected if they compared surgical interventions with conservative interventions or other surgical interventions in adults with fracture of the distal radius.

## DATA EXTRACTION

Data were extracted on patient characteristics, intervention, and outcomes (including functional and anatomic outcomes and complications). The quality of studies was assessed.

## MAIN RESULTS

44 studies (3193 mainly female and older patients with 3197 fractures) with 23 different comparisons met the selection criteria, with follow-up ranging from 6 weeks to 10 years. Summarizing the outcomes was impeded by the poor quality and variation in study methods, interventions, patient characteristics, and outcomes. Some anatomic and functional outcomes (Table) were better in the external-fixation group (7 studies), the pins-through-fracture group (4 studies), and the bone-scaffolding group (2 studies) than in the plaster-cast group; differences in func-

tion for open reduction and internal fixation or bone graft or substitute relative to plaster cast are not reported here because studies had excessive losses to follow-up or results were no longer significant when the random-effects model was used. External fixation and percutaneous pinning led to fewer patients with redisplacement that required secondary treatment than did a plaster cast; external fixation led to more patients with pin-track infection than did a plaster cast (Table). The evidence did not show clear superiority for 1 surgical intervention over another.

## CONCLUSIONS

In patients with distal radial fractures, heterogeneity exists for patients, mechanism of fracture, and fracture type, and results are inconsistent. Some benefit in reduced deformity, reduced malunion, and better functional outcome is seen for external fixation and percutaneous pinning relative to plaster cast, but who will benefit sufficiently is unclear.

*Sources of funding:* East Riding and Hull Health Authority, UK.

*For correspondence:* Professor R. Madhok, University of Hull, Willerby, England, UK. E-mail rajan.madhok@eriding-ha.northy.nhs.uk. ■

### Surgical interventions for fracture of the distal radius\*

Outcomes	Comparisons	Weighted event rates	RRR (95% CI)	NNT (CI)
Functional grading (FG) not excellent	Ext fix vs PC	55% vs 67%	18% (4 to 31)	9 (5 to 36)
	PTF vs PC	43% vs 81%	47% (29 to 61)	3 (2 to 5)
	Bone scaff vs PC	54% vs 78%	31% (13 to 45)	5 (3 to 10)
			RRI (CI)	NNH (CI)
	PSF vs PC	100% vs 84%†	19% (5.7 to 39)	7 (4 to 13)
			RRR (CI)	NNT (CI)
FG fair or poor	PTF vs PC	12% vs 37%	69% (36 to 85)	4 (3 to 9)
Redisplacement needing secondary treatment	Ext fix vs PC	0.9% vs 22%	89% (76 to 95)	5 (4 to 7)
	PF vs PC	0% vs 20%	92% (63 to 98)	6 (4 to 8)
			RRI (CI)	NNH (CI)
Pin track infection	Ext fix vs PC	11% vs 0%	648% (149 to 2148)	10 (7 to 16)

\*Bone scaff = bone scaffolding; Ext fix = external fixation; PC = plaster cast; PF = percutaneous fixation; PSF = pins supporting fracture; PTF = pins through fracture. Other abbreviations defined in Glossary; RRR, RRI, NNT, NNH, and CI calculated from data in article using a random-effects model.

†Event rates not weighted.

## COMMENTARY

Debate exists about the degree to which normal anatomy needs to be restored after distal radial fracture. Most clinicians agree that the articular surface should be made congruous, that shortening should be minimized, and that palmar tilt should be restored to at least neutral (1). The methods to achieve and maintain reduction can be grouped into casting and percutaneous or open techniques, although many variations of each exist.

Most clinicians consider high-energy multifragmentary distal radial fractures with intra-articular displacement as distinct from low-energy fractures with a congruous joint when choosing appropriate treatment. Many of the studies in the review by Handoll and Madhok have inappropriately evaluated all injuries together. The myriad of treatment 112 variations described, the wide range of outcomes evaluated, and the lack of injury discrimination have made it difficult to draw firm conclusions from the literature. Casting is associated with the highest probability of a poor anatomic result, especially in high-energy injuries and those with metaphyseal comminution. Alternatives to casting should be considered when acceptable reduction cannot be achieved or when

residual joint incongruity and metaphyseal redisplacement is likely, especially in high-energy injuries. It seems reasonable to approach these high-energy injuries with minimally invasive techniques and to resort to open reduction if anatomic restoration is not possible. The role of arthroscopy and adjunctive bone grafting or substitution to obtain and maintain reduction has yet to be elucidated (2).

Large prospective trials with appropriate injury discrimination and consistent outcome evaluation are needed before firm recommendations can be made about the degree of anatomic restoration required, and the best treatment to use in a particular situation. Existing practices are based largely on expert opinion.

*Hans J. Kreder, MD, MPH  
University of Toronto  
Toronto, Ontario, Canada*

## References

1. Hanel DP, Jones MD, Trumble TE. *Orthop Clin North Am.* 2002;33:35-57.
2. Trumble TE, Culp RW, Hanel DP, Geissler WB, Berger RA. *Instr Course Lect.* 1999;48:465-80.